## **CASE REPORT**

### FIXED DRUG ERUPTION

Balakrishna<sup>1</sup>, Souris Kondaveti<sup>2</sup>, Bhuvaneshwari<sup>3</sup>

#### **HOW TO CITE THIS ARTICLE:**

Balakrishna, Souris Kondaveti, Bhuvaneshwari. "Fixed Drug Eruption". Journal of Evidence based Medicine and Healthcare; Volume 2, Issue 7, February 16, 2015; Page: 936-938.

**ABSTRACT:** Fixed drug eruption is a drug induced skin eruption characterized by one or multiple skin erythematous plaques that disappears once the culprit drug is withdrawn leaving hyper pigmented sequelae. These plaques are termed fixed because they reappear on the same location each time the drug is administered. The commonly implicated drugs are tetracyclines, paracetamol and sulfonamides. A 38 yrs. old male patient presented to the department of medicine with history of erythematous patches all over the body. Patient was administered tablet Augmentin 625 mg for upper respiratory tract infection 4 days back. After taking the tablet he noticed a small erythematous patch on the right shoulder which increased in size and some patches are spread all over the body i.e. on back, proximal part of limbs and on abdomen. The patches were distinct sizes with hyperpigmentation and necrosis in the center. In the present case fixed drug eruption would be related to amoxicillin that is present in Augmentin in a view of a suggestive temporal relationship between drug intake and reaction onset, the remission of skin eruption after withdrawal. Hence caution is needed in patients who are hypersensitivity to amoxicillin.

**KEYWORDS:** Fix drug eruption, Augmentin.

**INTRODUCTION:** Fixed drug eruption is a drug induced skin eruption characterized by one or multiple skin erythematous plaques that disappears once the culprit drug is withdrawn leaving hyper pigmented sequelae.

**CASE DESCRIPTION:** A 38 yrs. old male patient presented to the department of medicine with history of erythematous patches all over the body. Patient was administered tablet augmentin for upper respiratory tract infection with phlegm four days back. After taking the tablet he noticed a small erythematous patch on the right shoulder which increased in size and later some patches are spread all over the body i.e. on back, proximal part of limbs and on abdomen. The patches were distinct sizes with hyperpigmentation and necrosis in the center. No history of fever, pain and burning sensation. No mucosal congestion or edema is seen. Vitals are within normal limits. No history of alcohol consumption, smoking and previous drug allergies. Patient was well built, conscious and coherent.

On general examination pulse, B.P, Respiratory rate and heart is within normal limits. Complete blood count, E.S.R, C.R.P and blood biochemistry including liver and renal function tests were normal.

Present case was diagnosed as fixed drug eruption and treated on outpatient basis with immediate stoppage of current medication and administration of oral anti-histamine levocetrizine 5mg.

## **CASE REPORT**

The case was reported to drug information center. Data regarding reaction was provided based on epidemiological studies and literature evidence. The information provided was that fixed drug eruption is a common reaction to amoxicillin that is present in Augmentin and also advised to perform patch test after 6 weeks to conclude the reaction. The reaction disappears in few days leaving hyper pigmented plaque. 6 weeks after complete healing of skin reaction, a patch test to Augmentin was performed both on the normal and involved skin simultaneously. Only patch test to Augmentin on involved skin was positive at 48hr due to reaction of residual hyperpigmentation.



Fig. 1

**DISCUSSION:** Present case diagnosed as fixed drug eruption to amoxicillin that is present in Augmentin. Narinjo and Uppsala monitoring scale used to assess the causality of drug reactions. The present reaction considered as PROBABLE as per WHO-UMC scale. The reaction is described moderate as per severity.

**CONCLUSION:** In the present case fixed drug eruption would be related to amoxicillin in a view of a suggestive temporal relationship between drug intake and reaction onset, the remission of skin eruption after withdrawal. Hence caution is needed in patients who are hypersensitivity to amoxicillin.

#### **REFERENCES:**

- 1. Sharma VK, Dhar S. Clinical pattern of cutaneous drug eruption among children and adolescents in north India. Pediatr Dermatol 1995; 12:178.
- 2. Mahboob A, Haroon TS. Drugs causing fixed eruptions: a study of 450 cases. Int J Dermatol 1998; 37:833.
- 3. Brahimi N, Routier E, Raison-Peyron N, et al. A three-year-analysis of fixed drug eruptions in hospital settings in France. Eur J Dermatol 2010; 20:461.
- 4. Nnoruka EN, Ikeh VO, Mbah AU. Fixed drug eruption in Nigeria. Int J Dermatol 2006; 45:1062.
- 5. Shiohara T, Mizukawa Y. Fixed drug eruption: a disease mediated by self-inflicted responses of intra epidermal T cells. Eur J Dermatol 2007; 17:201.
- 6. Shiohara T, Moriya N. Epidermal T cells: their functional role and disease relevance for dermatologists. J Invest Dermatol 1997; 109:271.

## **CASE REPORT**

- 7. Mizukawa Y, Yamazaki Y, Shiohara T. In vivo dynamics of intra epidermal CD8+ T cells and CD4+ T cells during the evolution of fixed drug eruption. Br J Dermatol 2008; 158:1230.
- 8. Shiohara T. Fixed drug eruption: pathogenesis and diagnostic tests. Curr Opin Allergy Clin Immunol 2009; 9:316.
- 9. Shiohara T, Mizukawa Y. The immunological basis of lichenoid tissue reaction. Autoimmun Rev 2005; 4:236.
- 10. Shelley WB, Shelley ED. Non pigmenting fixed drug eruption as a distinctive reaction pattern: examples caused by sensitivity to pseudoephedrine hydrochloride and tetrahydrozoline. J Am Acad Dermatol 1987; 17: 403.

#### **AUTHORS:**

- 1. Balakrishna
- 2. Souri S. Kondaveti
- 3. Bhuvaneshwari

#### PARTICULARS OF CONTRIBUTORS:

- 1. Junior Resident, Department of Pharmacology, Osmania General Hospital, Hyderabad.
- Assistant Professor, Department of Pharmacology, Osmania General Hospital, Hyderabad.

3. Assistant Professor, Department of Pharmacology, Osmania General Hospital, Hyderabad.

# NAME ADDRESS EMAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Souri S. Kondaveti,
Department of Pharmacology,
Osmania General Hospital, Hyderabad.
E-mail: drsourisuresh@gmail.com

Date of Submission: 09/02/2015. Date of Peer Review: 10/02/2015. Date of Acceptance: 11/02/2015. Date of Publishing: 14/02/2015.