

## ORIGINAL ARTICLE

### IDENTIFYING THE ROTATIONAL AXES OF DISTAL FEMUR IN SOUTH ANDHRA POPULATION OF INDIA USING IMAGE TOOL SOFTWARE

Sharmila Bhanu P<sup>1</sup>, Devi Sankar K<sup>2</sup>, Ravichandran D<sup>3</sup>, Ahammad Basha S<sup>4</sup>

#### HOW TO CITE THIS ARTICLE:

Sharmila Bhanu P, Devi Sankar K, Ravichandran D, Ahammad Basha S. "Identifying the Rotational Axes of Distal Femur in South Andhra Population of India Using Image Tool Software". Journal of Evidence based Medicine and Healthcare; Volume 2, Issue 32, August 10, 2015; Page: 4687-4694, DOI: 10.18410/jebmh/2015/659

**ABSTRACT:** **AIM:** To study the normal relationship of the anteroposterior (APA), transepicondylar (TEA) and posterior condylar (PCA) axis of normal cadaveric femoral bones using digital technology and special computer program. **MATERIAL AND METHOD:** The study comprised of 196 dry adult femora from 98 right and 98 left sides irrespective of sex and age belonging to Andhra Pradesh population of India. The bone collections were obtained from the Anatomy department, Narayana medical college, Nellore, India. The femurs were kept in normal anatomical position on OB. The photographs were taken from the distal end of all the femurs placing the camera lens 10cm constantly away from it with a digital camera. Using the reference points, angle between APA-TEA, APA-PCA and TEA-PCA were identified. The statistical significance of difference between the right and left groups was evaluated by using Student paired t-test. Data were presented as mean±SD. P-value less than 0.05 were considered statistically significant. **RESULTS:** The relationship of the angle between the APA-TEA, APA-PCA and TEA-PCA were observed. The angle of AP-TE, AP-PC and TE-PC was 94.84±3.43°, 87.64±1.62° and 6.84±2.71° respectively on right side. On the left side, the angle of AP-TE, AP-PC and TE-PC was 92.36±4.06°, 93.61±2.54° and 3.19±0.99° respectively. **CONCLUSION:** The normal femoral rotational alignment from cadaveric bone study using computer aided software can be helpful to the surgeon in selecting appropriate reference axis in any particular knee surgeries. In this regard the present data can be taken into consideration for the femoral rotational alignment during any intraoperative surgeries of knee and in total knee arthroplasty.

**KEYWORDS:** Femur, Rotational axes, Imaging software, Total knee arthroplasty, Distal femur.

**INTRODUCTION:** Accurate rotational alignment of the femoral component is an important surgical factor for successful total knee arthroplasty (TKA).<sup>1,2,3,4</sup> Improper rotation of the femoral implant during TKA may lead to abnormal patellofemoral kinematics, stiffness, abnormal gait patterns, flexion instability, asymmetrical flexion gaps and early failure.<sup>5,6,7</sup>

Various anatomic landmarks have been described for serving as rotational reference points for primary TKA. Three rotational axes referenced in distal femur for the placements of the femoral component during TKA are 1) anteroposterior axis (APA) or Whiteside's line, 2) transepicondylar axis (TEA) and 3) posterior condylar axis (PCA).<sup>8,9,10</sup>

Some degrees of external rotation of the femoral component are known to be crucial to prevent patellofemoral complications such as lateral tracking, subluxation and patellar component failure<sup>11</sup>. There is a controversy in the literature regarding the amount of external rotation and the ideal anatomical axis for optimal rotational alignment of the femoral component. Previous studies suggested that the APA of the distal femur was an easy and reliable landmark for

## ORIGINAL ARTICLE

rotational alignment of the femoral component, especially in the valgus knee.<sup>12,13,14</sup> However a number of studies have supported the clinical TEA i.e. the line connecting the medial and lateral epicondylar prominences, most consistently recreates a balanced flexion space and normal patello-femoral tracking.<sup>15</sup>

In the literature, most of the studies on these axes were in osteoarthritic knees during the arthroplasty procedures, the results of which might not represent normal relationship due to the deformed bones in those cases. Moreover reference axis from plain X-ray, CT or MRI scan can be inaccurate because most of these will be diseased cases of the knee and the axes measured may not be as accurate as in the normal subjects.<sup>13,16</sup>

Knowing the normal relationship of the above said three axes, it can be helpful to the surgeon in selecting one or more appropriate reference axis to properly rotate the femoral component in any particular knee during the operation.<sup>17,18</sup> In the present study computer software is applied to study the relationship among the APA, PCA and TEA in cadaveric femurs instead of using manual goniometer.

**MATERIAL AND METHODS:** The study comprised of 196 dry adult femora from 98 right and 98 left sides irrespective of sex and age belonging to Andhra Pradesh population of India. The bone collection was obtained from the Anatomy department, Narayana medical college, Nellore, India.

As criteria of inclusion, none of the femora presented fractures, malformations, damage due to conservation or pathologies that could influence the development of the studied region. Each of the femurs was digitally photographed in standardized position on acrylic osteometric board (OB). The femurs were kept on normal anatomical position, of which the posterior most point of medial and lateral condyles (distally) i.e. posterior condylar axis (PCA), and the greater trochanter (proximally) were rested on OB and the anterior surface of shaft directed upwards. The photographs were taken from the inferior surface of distal end of all the femurs by placing the camera lens 10cm constantly away from it (Fig. 1) with the help of a digital camera (Sony DCR W270, Tokyo, Japan). The stored images were transferred to a PC unit.<sup>19</sup>

All the digital images were processed to locate the six reference points and three lines using Adobe Photoshop 7.0 for Windows and the final images were saved in jpeg format (Fig. 2a). The formatted images were transferred to Image Tool Software (UTHSCSA Image Tool for Windows version 3.0, San Antonio, TX, US) and analyzed (Fig. 2b). With the help of reference points the angle between APA-TEA, APA-PCA and TEA-PCA were identified. The data was fed in computer program SPSS ver.10 for Windows (SPSS Inc., Chicago, IL, USA). The statistical significance of difference between the right and left groups was evaluated by using Student paired t-test. Data were presented as mean±SD. P-value less than 0.05 were considered statistically significant.

### **Axes measured in the distal end of femur were,**

1. APA: Anteroposterior axis is the line drawn from the trochlear or patellar groove, anteriorly to the apex or center of the intercondylar notch, posteriorly.<sup>20</sup>
2. TEA: Trans-epicondylar axis is the line joining the most prominent point of the medial and lateral epicondyles of femur.<sup>21</sup>
3. PCA: Posterior condylar axis is the tangent line joining the posterior point of the medial and lateral femoral condyles.<sup>7</sup>

## ORIGINAL ARTICLE

---

Angulations calculated between these three axes were AP-TE; AP-PC and TE-PC (Fig.2b)

**RESULTS:** The relationship of the angle between the APA-TEA, APA-PCA and TEA-PCA were observed. The angle of AP-TE, AP-PC and TE-PC was  $94.84 \pm 3.43^\circ$ ,  $87.64 \pm 1.62^\circ$  and  $6.84 \pm 2.71^\circ$  respectively on right side. On the left side, the angle of AP-TE, AP-PC and TE-PC was  $92.36 \pm 4.06^\circ$ ,  $93.61 \pm 2.54^\circ$  and  $3.19 \pm 0.99^\circ$  respectively (Table 1).

**DISCUSSION:** There is no compromise regarding the use of the axes and angles for measuring the rotation of the femoral component. Rotation arrangement of the femoral component in TKA is imperative for the outcome of the surgery. Malrotation of the femoral component on the femur may lead to patellofemoral dislocation or subluxation, to wear or loosening of the patellar component.<sup>22</sup> The correlation of AP-TE; AP-PC and TE-PC axes and angles noted in the literature vary as they are found to have different methodology in each study.

Accomplishing optimal femoral component rotational alignment in TKA is crucial in establishing a balanced knee reconstruction and ensuring adequate patello-femoral tracking. Unbalanced knees can lead to instability, patellofemoral problems, persistent pain, stiffness, and generally poorer outcomes including early failure.<sup>23</sup> Although the desired positions and guiding landmarks for placement of the femoral and tibial components in the coronal and sagittal planes have been well described, rotational positioning of the components can still be problematic.<sup>24,25,26</sup> Performing a TKA requires the accurate execution of key bone cuts in the correct orientation to the appropriate axes. There is huge potential for cumulative errors to occur, which may have significant and dramatic effects on function and longevity.<sup>16</sup>

A number of studies are found in relation to the anatomy and functional axes of the normal femur.<sup>13,27,28,29</sup> These include the use of the TEA, APA and PCA. Various intraoperative techniques have been described to achieve the optimal femoral component rotation. The axis of the posterior femoral condyles has normally been used as the reference for neutral rotation of the femur.<sup>30</sup> This was thought to be a reasonable landmark because the goal of TKA is to re-establish alignment of the anatomic femoral condyles.<sup>11,31</sup> However, recent biomechanical analyses using instant centers of motion analysis have suggested that the TEA parallels the primary center of rotation of the knee joint but not the posterior condylar axis.<sup>23</sup> The fixed axis that closely approximate the epicondylar axis reported is the TEA has currently attracted attention as an ideal rotational reference that provides functional kinematics (31 Doro LC). Furthermore, several authors also reported that the TEA most consistently recreates a balanced flexion gap.<sup>3</sup>

In a study on 100 arthritic knees in TKA reported by 32 Poilvache et al, the AP-TE, AP-PC and TE-PC angle was  $90.33 \pm 2.44$ ,  $86.92 \pm 2.71$  and  $3.60 \pm 2.02$  degrees, respectively. Matsuda et al<sup>3</sup> findings of the femoral measurements showed the AP-PC and TE-PC angles of  $83.70 \pm 2.44$ ,  $6.03 \pm 3.60$  degrees in normal knees and  $83.43 \pm 2.54$  and  $6.00 \pm 2.35$  degrees in varus knees, in which AP-TE angle was not reported. Arima et al<sup>12</sup> studied AP-PC and TE-PC angles in normal femurs by visual measurement were  $86.2 \pm 2.0$  and  $4.4 \pm 2.9$  degrees and by radiographic measurement were  $86.9 \pm 1.7$  and  $5.7 \pm 1.7$  degrees respectively, with radiographic measurement less than the visual measurement.

## ORIGINAL ARTICLE

---

In the present study relationship of the angle between the AP-TE, AP-PC and TE-PC was  $94.84\pm 3.43^\circ$ ,  $87.64\pm 1.62^\circ$  and  $6.84\pm 2.71^\circ$  respectively on right side. On the left side, the angle of AP-TE, AP-PC and TE-PC was  $92.36\pm 4.06^\circ$ ,  $93.61\pm 2.54^\circ$  and  $3.19\pm 0.99^\circ$  respectively. These observations were almost coinciding with the other findings and is found to significant ( $p<0.0001$ ). The comparison of angles of all the femurs observed between AP-TE and AP-PC was  $93.60\pm 3.95$  and  $90.62\pm 3.67$  respectively with the mean difference  $2.98\pm 5.18$  was also found to be significant (Table 2). However this study is only restricted to the normal femurs but not to the arthritic and other deformed femoral axes which may differ from the present study.

Using image tool software technique, the relations between AP, TE and PC axes in normal dry femora observed in the present study may provide a minimal error of these measurements, as these measurements are mostly done on the surgical table at the time of TKA procedures. The clinical TEA is found to be more significant than the other axis in this study as well with the other studies. As this TEA has the perpendicularity to the mechanical axis of femur and tibia with knee fixed at  $90^\circ$ ,<sup>33,34</sup> makes it a sound landmark for rotational alignment of femurs in TKA.

Even though the epicondyles have been considered as the reliable anatomic landmarks, it is difficult to impound its peak exactly during the surgery.<sup>12</sup> Since the aim of TKA is to achieve a normal rotational alignment of the femur, a multi reference axes should be used rather depending upon any single rotational axis of the distal femur.<sup>35</sup> Though there are many controversial debates regarding the rotational measurements, the normal cadaveric bone study in particularly the computer aided image software methods can be relied upon than the visual calculated measurements. In this context, the normal anatomical knowledge of rotational axes of distal femur namely AP, PC and TE and its association in the cadaveric dry bones must be studied and verified. The data presented in this study therefore may be taken into deliberation in the analysis of the rotational axes during TKA.

### REFERENCES:

1. Akagi M, Yamashita E, Nakagawa T, Asano T, Nakamura T: Relationship between frontal knee alignment and reference axes in the distal femur. *Clin Orthop Relat Res*, 2001; 388: 147-56.
2. Barrack RL, Schrader T, Bertot AJ, Wolfe MW, Myers L: Component rotation and anterior knee pain after total knee arthroplasty. *Clin Orthop Relat Res*, 392: 46-55, 2001.
3. Matsuda S, Miura H, Nagamine R, Urabe K, Mawatari T, Iwamoto Y: A comparison of rotational landmarks in the distal femur and the tibial shaft. *Clin Orthop Relat Res*, 2003; 414: 183-88.
4. Miller MC, Berger RA, Petrella AJ, Karmas A, Rubash HE. Optimizing femoral component rotation in total knee arthroplasty. *Clin Orthop Relat Res* 2001; 392: 38-45.
5. Victor J. Rotational alignment of the distal femur: A literature review. *Orthop Traumatol Surg Res* 2009; 95: 365-72.
6. Aglietti P, Sensi L, Cuomo P, Ciardullo A. Rotational position of femoral and tibial components in TKA using the femoral transepicondylar axis. *Clin Orthop Relat Res* 2008; 466: 2751-55.

## ORIGINAL ARTICLE

---

7. Rossi R, Bruzzone M, Bonasia DE, Marmotti A, Castoldi F. Evaluation of tibial rotational alignment in total knee arthroplasty: a cadaver study. *Knee Surg Sports Traumatol Arthrosc* 2010; 18: 889-93.
8. Mullaji AB, Sharma AK, Marawar SV, Kohli AF, Singh DP. Distal femoral rotational axes in Indian knees. *J Orthop Surg*. 2009; 17: 166-69.
9. Olcott CW, Scott RD. A comparison of 4 intraoperative methods to determine femoral component rotation during total knee arthroplasty. *J Arthroplasty* 2000; 15: 22-6.
10. Berhouet J, Beaufils P, Boisrenoult P, Frasca D, Pujol N. Rotational positioning of the tibial tray in total knee arthroplasty: a CT evaluation. *Orthop Traumatol Surg Res* 2011; 97: 699-704.
11. Zhang XL, Zhang W, Shao JJ. Rotational alignment in total knee arthroplasty: nonimage-based navigation system versus conventional technique. *Chin Med J* 2012; 125(2): 236-243.
12. Arima J, Whiteside LA, McCarthy DS, White SE. Femoral rotational alignment, based on the anteroposterior axis, in total knee arthroplasty in a valgus knee. A technical note. *J Bone Joint Surg Am* 1995; 77: 1331-4.
13. Saengnipanthkul S, Sarntipiphath C, Chaichoon A, Chaisiwamongkol G. The relationship of anteroposterior, transepicondylar and posterior condylar axis of the femur: A Thai cadaveric study using digital measurement. *J Med Assoc Thai* 2009; 92 (10): 1295-9.
14. Classen T, Wegner A, Muller RD, Von Knoch M. Femoral component rotation and Laurin angle after total knee arthroplasty. *Acta Orthop Belg* 2010; 76: 69-73.
15. Insall JN, Scuderi GR, Komistek RD, Math K, Dennis DA, Anderson DT. Correlation between condylar lift-off and femoral component alignment. *Clin Orthop Relat Res* 2002; 403: 143-52.
16. Luo CF. Reference axes for reconstruction of the knee. *Knee* 2004; 11: 251-57.
17. Kinzel V, Ledger M, Shakespeare D. Can the epicondylar axis be defined accurately in total knee arthroplasty? *Knee* 2005; 12: 293-96.
18. Middleton FR, Palmer SH. How accurate is Whiteside's line as a reference axis in total knee arthroplasty? *Knee* 2007; 14: 204-07.
19. Ravichandran D, Devi Sankar K, Sharmila Bhanu P, Manjunath KY, Shankar R. Angle of femoral neck anteversion in Andhra Pradesh population of India using image tool software. *JIMSA* 2014; 27: 199-200.
20. Chang CB, Seong SC, Lee S, Lee MC. Anatomical assessment of the distal femur and tibia for optimal femoral rotational alignment in total knee arthroplasty. *J Korean Knee Soc*. 2010; 22: 46-54.
21. Aglietti P, Sensi L, Cuomo P, Ciardullo A. Rotational position of femoral and tibial components in TKA using the femoral transepicondylar axis. *Clin Orthop Relat Res* 2008; 466: 2751-755.
22. Amiss AA, Firer P, Mountney J, Senavongse W, Thomas NP. Anatomy and biomechanics of the medial patellofemoral ligament. *Knee*, 2003; 10: 215-20.
23. Vaidya SV, Gadhiya RM, Bagaria V, Ranawat AS, Ranawat CS. Computed tomographic evaluation of femoral component rotation in total knee arthroplasty. *Indian J Orthop*. 2013; 47:40-4.

## ORIGINAL ARTICLE

24. Goodman SB, Patel JJ, Delp SL, Giori NJ. The high variability of tibial rotational alignment in total knee arthroplasty. *Clin Orthop* 2006; 452: 65-69.
25. Barrack RL, Schrader T, Bertot AJ, Wolfe MW, Myers L: Component rotation and anterior knee pain after total knee arthroplasty. *Clin Orthop Relat Res*, 392: 46-55, 2001.
26. Chang CB, Seong SC, Lee S, Yoo JH, Rhee SH, Lee MC: Anatomical assessment of distal femur for optimal femoral component rotational alignment in TKA. *J Korean Orthop Assoc*, 40: 882-888, 2005.
27. Lie WHC, Chiu KY, Ng TP. Radiological assessment of femoral rotation: A cadaveric study. *J Orthop Trau and Rehab* 2012; 16: 22-25.
28. Tang WM, Chiu KY, Kwan MF, Ng TP, Yau WP. Sagittal bowing of the distal femur in Chinese patients who require total knee arthroplasty. *J Orthop Res* 2005; 23: 41-5.
29. Yau WP, Chiu KY, Tang WM, Ng TP. Coronal bowing of the femur and tibia in Chinese: its incidence and effects on total knee arthroplasty planning. *J Orthop Surg (Hong Kong)* 2007; 15: 32-6.
30. Ghosh KM, Merican AM, Iranpour F, Deehan DJ, Amis AA. The Effect of Femoral Component Rotation on the Extensor Retinaculum of the Knee. *J Orthop Res* 2010; 28: 1136-41.
31. Siston RA, Patel JJ, Goodman SB, Delp SL, Giori NJ. The Variability of femoral rotational alignment in total knee arthroplasty. *J Bone Joint Surg Am.* 2005; 87: 2276-80.
32. Doro LC, Hughes RE, Miller JD, Schultz KF, Hallstrom B, Urquhart AG. The reproducibility of a kinematically-derived axis of the knee versus digitized anatomical landmarks using a knee navigation system. *The Open Biomedical Engineering Journal*, 2008, 2, 52-56.
33. Poilvache PL, Insall JN, Scuderi GR, Font-Rodriguez DE. Rotational landmarks and sizing of the distal femur in total knee arthroplasty. *Clin Orthop Relat Res* 1996; 331: 35-46.
34. Stiehl JB, Cherveny PM. Femoral rotational alignment using the tibial shaft axis in total knee arthroplasty. *Clin Orthop* 1996; (331): 47-55.
35. Yau WP, Leung A, Liu KG, Yan CH, Wong LS, Chiu KY. Errors in the identification of the transepicondylar and anteroposterior axes of the distal femur in total knee replacement using minimally-invasive and conventional approaches: a cadaver study. *J Bone Joint Surg Br* 2008; 90: 520-6.

<b>Variables</b>	<b>Right side Mean±SD (In degrees) n=98</b>	<b>Left side Mean±SD (In degrees) n=98</b>	<b>t-value</b>	<b>p-value</b>
AP-TE	94.84±3.43	92.36±4.07	4.62	< 0.0001*
AP-PC	87.64±1.62	93.61±2.54	19.61	< 0.0001*
TE-PC	6.84±2.71	3.19±0.99	19.61	< 0.0001*

**Table 1: Comparison between right and left femurs for AP-TE, AP-TC and TE-PC**

Independent sample T-Test was applied. \* represents extreme significant.


# ORIGINAL ARTICLE

FEMUR		Mean±SD (In degrees) n=196	Paired Difference	t-value	p-value
			Mean±SD (In degrees)		
Pair 1	AP-TE	93.60±3.95	2.98±5.18	8.048	< 0.0001*
	AP-PC	90.62±3.67			
Pair 2	AP-TE	93.60±3.95	88.58±3.19	389.130	< 0.0001*
	TE-PC	5.01±2.74			
Pair 3	AP-PC	90.63±3.67	85.61±5.54	216.192	< 0.0001*
	TE-PC	5.02±2.74			

Table 2: Comparison of both side of femurs for AP-TE, AP-TC and TE-PC

Paired sample T-Test was applied. \* represents extreme significant.

**Figure 1:** The orientation of femur and camera on the osteometric board. A-optical axis; B-anatomical axis; C-camera lens.



**Figure 1**

**Figure 2a:** Schematic diagram of distal femur showing the rotational axes and its angles. APA–anteroposterior axis; TEA–transepicondylar axis; PCA–posterior condylar axis; AP-TE–angle between APA and TEA; AP-PC–angle between APA and PCA; TEA-PCA–angle between TEA and PCA; TEA’–parallel line to TEA; 1-6–reference points (Red dots). **2b:** Screenshot image of distal femur and its axes for measurement of angles using image tool software in Windows.




Figure 2a &amp; 2b

**AUTHORS:**

1. Sharmila Bhanu P.
2. Devi Sankar K.
3. Ravichandran D.
4. Ahammad Basha S.

**PARTICULARS OF CONTRIBUTORS:**

1. Assistant Professor, Department of Anatomy, Narayana Medical College & GH, Nellore, Andhra Pradesh, India.
2. Assistant Professor, Department of Anatomy, Narayana Medical College & GH, Nellore, Andhra Pradesh, India.
3. Professor, Department of Anatomy, Karpagam Faculty of Medical Sciences & Research, Coimbatore, Tamil Nadu, India.

4. Assistant Professor, Department of Social & Preventive Medicine, Narayana Medical College & GH, Nellore, Andhra Pradesh.

**NAME ADDRESS EMAIL ID OF THE CORRESPONDING AUTHOR:**

Dr. Sharmila Bhanu P,  
Assistant Professor,  
Department of Anatomy,  
Narayana Medical College & GH,  
Nellore-524003, Andhra Pradesh, India.  
E-mail: sharmilabp78@gmail.com

Date of Submission: 22/07/2015.  
Date of Peer Review: 23/07/2015.  
Date of Acceptance: 28/07/2015.  
Date of Publishing: 04/08/2015.