

## CASE REPORT

### ACUTE BILATERAL VIRAL NECROTIZING RETINITIS: AN UNCOMMON CASE REPORT

Rajendra Ku. Behera<sup>1</sup>, R. C. Mahapatro<sup>2</sup>, J. P. Behera<sup>3</sup>, Altawab Bassel<sup>4</sup>, M. K. Bal<sup>5</sup>

#### HOW TO CITE THIS ARTICLE:

Rajendra Ku. Behera, R. C. Mahapatro, J. P. Behera, Altawab Bassel, M. K. Bal. "Acute Bilateral Viral Necrotizing Retinitis: An Uncommon Case Report". Journal of Evidence based Medicine and Healthcare; Volume 2, Issue 35, August 31, 2015; Page: 5551-5554, DOI: 10.18410/jebmh/2015/769

**ABSTRACT:** A 22 year old male with a history of high grade fever 2 days, diarrhea 3 times and vomiting 2 times presented with diminution of vision in right eye of 1 days duration. His best corrected visual acuity (BCVA) was counting finger 1 meter with no pin hole improvement and 20/20 (Snellen's) in the right and left eye respectively. Fundus examination RE revealed white lesion in geographic fashion with clear edge involving macula and in left eye small peanut size white lesion present at paramacular area. Clinically a diagnosis of acute necrotizing was made. We started treatment by intra venous antiviral and systemic steroid. ELISA (serum) and PCR (aqueous) were positive for herpes simplex virus (Index above 1.1 i.e. 1.54).<sup>1,2</sup> The lesions showed a good response to the above treatment. At 2 months follow-up, lesion had resolved well with BCVA of 20/40 and 20/20 in right and left eye respectively.

**KEYWORDS:** Bilateral, Herpes simplex, Retinitis.


**INTRODUCTION:** Herpes viruses are a DNA virus that affect man and animals & cause multiple systemic diseases. Herpes simplex occurs naturally only in man. Ophthalmic manifestation includes keratitis, follicular conjunctivitis, choreoretinitis and uncommonly acute necrotizing retinitis, which is a serious manifestation.<sup>3,4</sup> We report an interesting case of bilateral necrotizing retinitis and visual prognosis with close follow up.

**CASE REPORT:** A 22 years old male (Student) presented with sudden painless loss of vision in the right eye of 1 day duration. Patient was symptomatically treated for fever, diarrhea and vomiting for 2 days. On examination his BCVA was CF 1 meter and 20/20 in the right and left eye respectively. Severe RAPD was noted in right eye. Slit lamp examination showed a quiet anterior segment and 1+ vitreous cells (Grading: Himura and Colleagues) in both eye, IOP were normal. Fundus examination of right eye showed area of necrotizing retinitis in geographic fashion involving macula. In the left eye a peanut size white patches were seen temporal to macula (Fig. 1 A & B). Fundus fluorescein angiography showed area of capillary non-perfusion corresponding to retinitis lesion and optical coherence tomography showed area of retinal full thickness necrosis. Laboratory investigation includes an ESR-46mm 1<sup>st</sup> hour, Hb%-12.7 mg%, TWBC-10,800/Cumm, DC-N 75%, L-15%, E-10%, urine contain trace albumin. Coagulation profile, blood sugar, liver and renal function, chest X-ray normal. VDRL (slide flocculation) are negative. ELISA showed IgM and IgG antibodies positive for herpes simplex I and II. Aqueous tap was +ve for herpes simplex virus.<sup>5</sup>

He was treated with intravenous 750mg valacyclovir (1500mg/m<sup>2</sup>/day) first three days, then next oral steroid in a dose of 1mg/kg body weight were added on the second day of antiviral treatment. Then antiviral 1000mg tab BID for 7 days and 500 mg tab BID for next 7 days and

## CASE REPORT


steroid gradually taper over 4 weeks.<sup>6</sup> At the end of 4 weeks there was no further deterioration in vision or appearance of any new lesions. At this last follow up (1 month) the retinal lesions had healed well (Figure-1a and 1b) and has BCVA was 20/40 and 20/20 in the right and left eyes respectively.



**Fig. 1a & 1b: Photograph showing the 1<sup>st</sup> day**

**DISCUSSION:** The most common clinical manifestation of HSV retinitis has been described as acute retinal necrosis (ARN) syndrome in the Japanese literature by Urayama et al in 1971.<sup>7</sup> But in 1990 Foster et al described the entity as progressive acute retinal necrosis (PORN) in AIDS patients having severe immunocompromised state.<sup>8</sup> We report an interesting case of bilateral retinal necrosis in a patient with PCR and aqueous sample proved positive for Herpes simplex virus. It is an adult male suffered from sudden diminution of vision following a bout of high grade fever. So though bilateral involvement may be due to viremic spread rather than neural spread in the most consistent key feature.<sup>9</sup>

Most cases are associated with encephalitis, but absent in this case. Also typical history of ocular pain, mild anterior uveitis is absent except mild vitritis. Retinal necrosis results in rapid confluence of well-demarcated areas of full thickness retinal involvement over the next 2 to 4 days may be due to an occlusive vasculitis and papillitis.<sup>10</sup> Basing on clinical diagnosis, we started I.V. antiviral and then systemic steroid to prevent further retinal damage and got good visual improvement as well as complete regression of the lesion which was remarkable.



**Fig. 2a & 2b: Photograph showing the 9<sup>th</sup> day**

## CASE REPORT

---

### RECORD OF THE VA (RIGHT EYE):

- 1 Day - 1 meter.
- 3 Days - 4 meter.
- 7 Days - 20/100.
- 9 Days - 20/80.
- 2 Weeks - 20/70.
- 4 Weeks - 20/40.
- 8 Weeks - 20/40.

### REFERENCES:

1. Madhavan HN, Priya K. The diagnostic significance of ALISA for herpes simplex, varicella zoster and CMV retinitis. *Indian J Ophthalmology*, 2003; 51: 71-5.
2. Grover R, Ratho R.K. et al: Role of viral serology in the diagnosis of acute retinal necrosis syndrome. *Indian J. Patholmicrobiol*, 2002; 15: 269-71.
3. RA, C.K J Paniker. Text book of microbiology, 4<sup>th</sup> edn page 461-469.
4. Culbertson WW, et al: Varicella Zoster Virus ~ a cause of the acute retinal necrosis syndrome. *Ophthalmology*, 1986; 93: 559-69.
5. Holland GN et al: An association between acute retinal necrosis syndrome and HLA types. *Am. J. Ophthalmology*, 1989; 108 (4): 370-4.
6. Blumenkranz MS, et al: Treatment of the acute retinal necrosis syndrome with intravenous acyclovir. *Ophthalmology*, 1986; 93: 296-300.
7. Urayama A., Yamada N, Sasaki T, et al: Unilateral acute uveitis with retinal periarthritis and retinal detachment. *Jpn. J. Clin. Ophthalmol.*, 1971; 25: 607-29.
8. Forster DJ, Rao NA et al: Rapidly progressive outer retinal necrosis in the AIDS. *Am. J. Ophthalmol.* 1990; 110: 391-8.
9. Duker JS, Nielsen JC et: Rapidly progressive acute retinal necrosis secondary to herpes simplex virus type-1. *Ophthalmology*, 1990; 97: 1638-43.
10. Myran Yanoff, Jay S. Dukar. Text Book of Ophthalmology, Page 10.11.4 to 10.11.6.

## CASE REPORT

---

### **AUTHORS:**

1. Rajendra Ku. Behera
2. R. C. Mahapatro
3. J. P. Behera
4. Altawab Bassel
5. M. K. Bal

### **PARTICULARS OF CONTRIBUTORS:**

1. Assistant Professor, Department of Ophthalmology, Advances Base Care Clinic.
2. Professor, Department of Ophthalmology, Advances Base Care Clinic.
3. Professor, Department of Ophthalmology, Advances Base Care Clinic.

4. Assistant Professor, Department of Ophthalmology, Advances Base Care Clinic.
5. Director, Department of Ophthalmology, Advances Base Care Clinic.

### **NAME ADDRESS EMAIL ID OF THE CORRESPONDING AUTHOR:**

Dr. Rajendra Ku. Behera,  
M. K. C. G. Medical College,  
Qtr. 3R/23. Odisha-760001.  
E-mail: drrajendra04@yahoo.com

Date of Submission: 17/08/2015.  
Date of Peer Review: 18/08/2015.  
Date of Acceptance: 20/08/2015.  
Date of Publishing: 31/08/2015.